

14. Поляков, И.Я. Прогноз появления и учет вредителей и болезней сельскохозяйственных культур [Текст] / И.Я.Поляков. – М.: МСХ СССР, 1958. – 632 с.
15. Строгая, Г.М. Биология яблонной моли Южной Киргизии и использование биологического метода в борьбе с ней [Текст] / Г.М.Строгая // В тр. Плодовые леса Южной Киргизии и использование.- М-Л.: АН СССР, 1949.
16. Тешебаева, З.А. Энтомопатогенные микроорганизмы непарного шелкопряда (*Lymantria dispar* L.) в орехово-плодовых лесах Кыргызстана [Текст]: автореф. дис. ... канд. биол. наук. / З.А.Тешебаева.- Бишкек, 2012. - 26с.
17. Токторалиев, Б.А. Защита орехово-плодовых лесов Кыргызстана от насекомых- вредителей [Текст] / [Б.А.Токторалиев, З.А. Тешебаева, Дей Жун Ш., Чан Х., Чан Ли Ч., А.Т. Аттокуров] // Известия Ошского технологического университета. – Ош: ОшТУ, 2018.- № 2.- С. 150-155.
18. Токторалиев, А.А. Биология и экология горного кольчатого шелкопряда в условиях орехоплодовых лесов южного Кыргызстана [Текст]: автореф. дис. ... канд. биол. наук. / А.А.Токторалиев. - Бишкек, 2003. - 13с.
19. Токторалиев, Б.А. Экологизация методов защиты леса в Кыргызстане [Текст] / З.А.Тешебаева, В.Ш.Исабекова // Известия Кыргызского государственного технического университета им. И. Раззакова.- Бишкек, 2018.- №3.- С.342-348.
20. Фасулати, К.К. Полевое изучение наземных беспозвоночных [Текст] / К.К.Фасулати. - М.: Высшая школа, 1971. – 424 с.
21. Ходырев, В.П. Энтомопатогенные микроорганизмы в очагах непарного шелкопряда (*Lymantria dispar* L.) орехово-плодовых лесов юга Кыргызстана [Текст] / [В.П.Ходырев, С.А. Бахвалов, З.А.Тешебаева, Б.А.Токторалиев] // Сибирский экологический журнал. - 2010. - Т. 17. - № 5. - С. 701-707.
22. Самиева, Ж.Т. Современные пути решения проблемы повышения рентабельности и экологизации сельскохозяйственного производства и его переработки [Текст] / Ж.Т. Самиева // Наука. Образование. Техника. – Ош: КУУ, 2019. – №1. – С. 122 - 129.

DOI: 10. 54834 / 16945220_2021_3_58

Поступила в редакцию 15. 09. 2021 г.

УДК 634.582.232.

Сагитов А.О.

академик НАН Республики Казахстан

Калыкова Г.Н.

науч. сотруд. научно-произв. цент. Института Биологии НАН Кыргызской Республики

Тешебаева З.А.

к.б.н, нач. отд. науки Ошского технол. универ. им. М.М. Адышева, Кыргызская Республика

Кунсуралиева И.К.

к.б.н., науч. сотруд. научно-произв. цент. инст. биологии НАН Кыргызской Республики

КЫРГЫЗСТАНДА СЕМЕНОВ ПИХТАСЫНЫН ИЛДЕТТЕРИ ЖАНА ЗЫЯНКЕЧТЕРИ МЕНЕН КҮРӨШҮҮ ЧАРАЛАРЫ

Батыш Тянь Шандагы Семенов ак карагай (Abies. Semonovii Fedtsch, 1898) токойлорунун санитардык абалына баа берүү изилдөөнүн предмети болуп эсептелет. Изилдөөнүн максаты ак карагай токойлорунун илдеттери менен зыянкечтерин тактоо жана аларга каршы күрөшү иш аракеттеринин натыйжалуулугун көрсөтүү болду. Карагайлуу токойлордо илдет чакыруучу козу карындар менен зыянкечтердин саны кескин өсүп, санитардык абалы канааттандырарлык эмес деп баалоого болот. Жалпылап караганда токойдо энтомологиялык жана фитопатологиялык зыянкечтердин болуусу жаратылыш биоартүрдүүлүгүнүн бир бөлүгү болуп каралганы менен, мындай процесстердин улануусунан дарактардын соолуп куурап калышынан табыйгый өсүп таралуусун токтоткон. Мындай көрүнүшкө себепкер пихта токойюнда кеңири тараган

зыянкечтер катары тобурчак зыянкечи (*Dioryctria abietella* Schiff.1775), уруктун зыянкечтери *Megastigmus Dalm.*, карагай пихта хермеси (*Aphrastasia pectinatae* Chol.1888). Ак карагайдын карындар катары (*Trichothecium roseum*(Pers.) Link, 1809) жана (*Myxotrichum chartarum*, Kunze, 1823).

Негизги сөздөр: ак карагай; ак карагай токойлору; козу карын; кунөскана илдет чакыруучу козу карындар; өнүм; зыянкечтер; фунгицид; биоартурдүүлүк; илдетти; өнүм.

БОЛЕЗНИ И ВРЕДИТЕЛИ ПИХТЫ СЕМЕНОВА В КЫРГЫЗСТАНЕ И МЕРЫ БОРЬБЫ С НИМИ

Предметом исследования является оценка санитарного состояния пихтового леса (*Abies. Semenovii* Fedtsch, 1898) Западного Тянь-Шаня. Целью исследования является выявление болезней и вредителей пихтовых лесов и демонстрация эффективности мер борьбы. Санитарному состоянию пихтовых лесов в настоящее время можно дать относительно положительную оценку. Тем не менее, в них встречаются вредители и грибные заболевания. В целом наличие в лесу энтомологических и фитопатологических вредителей, это тоже часть всего лесного биоразнообразия, созданного самой природой. Санитарному состоянию пихтовых лесов в настоящее время можно дать относительно положительную оценку. Тем не менее, в них встречаются вредные организмы. В процессе жизнедеятельности этих вредных организмов здоровые деревья усыхают, насаждения теряют свою роль и способность к естественному возобновлению. Наиболее распространенными вредителями пихтовых лесов являются шишковая огневка (*Dioryctria abietella* Schiff.1775), семяеды рода *Megastigmus* Dalm., хермес елово-пихтовый бурый (*Aphrastasia pectinatae* Chol.1888). Среди болезней пихты Семенова наиболее широкое распространение в питомнике имеет фузариозное поражение всходов, а в естественных насаждениях – ржавчина (*Melampsorella cerastii* Wint.), и ценангиальный рак (*Cenangium abietis* (Pers.) Reh. Duby, 1896). Другое наиболее распространенное заболевание – альтернариоз, вызываемое несовершенным грибом вида (*Alternaria alternata* (Fr.) Keissle, 1912). Наиболее опасное заболевание пихты Семенова корневая гниль, вызываемая корневой губкой (*Fomitopsis annosa* (Fr.) Karst.1881). Также на пихте Семенова обнаружены, возбудители опасных грибных заболеваний (*Trichothecium roseum*(Pers.) Link, 1809) и (*Myxotrichum chartarum*, Kunze, 1823).

Ключевые слова: пихта; пихтовые леса; грибы; питомник; возбудители болезней; вредители; фунгициды; биоразнообразие; заболевания; всходы.

PESTS AND DISEASES OF SEMENOVII FIR IN KYRGYZSTAN AND MEASURES TO COMBAT THEM

The subject of the study was the assessment of the sanitary state of fir forest in the western Tien Shan. The aim of the study was to identify diseases and pests of fir forests and to demonstrate the effectiveness of control measures. The sanitary condition of fir forests can currently be assessed relatively positively. Nevertheless, hazardous organisms are found in them. In the process of vital activity of these hazardous organisms, healthy trees dry out, plantings lose their role and ability to natural regeneration. Abstract. In general, the presence of entomological and phytopathological pests in the forest is also part of the entire forest biodiversity created by nature itself. The sanitary condition of fir forests can currently be assessed relatively positive. Nevertheless, hazardous organisms are found in them. In the process of vital activity of these hazardous organisms, healthy trees dry out, plantings lose their role and ability to natural regeneration. The most common pests of fir forests are pine *Dioryctria abietella* Schiff., *Megastigmus* (*Megastigmus* Sp.), *Adelges* (*Aphrastasia pectinatae* Chol.) Among the diseases of Semyonov fir, the most widespread in the nursery is fusarium lesion of seedlings, and in natural stands – rust (*Melampsorella cerastii* Wint.), and cenangium cancer (*Cenangium abietis* (Pers.) Reh. Duby, 1896). Another most common disease is *Alternaria* blight caused by an imperfect fungus *Alternaria alternata* (Fr.) Keissler. The most dangerous disease of Semyonov fir is root rot, caused by the pine fungus *Fomitopsis annosa* (Fr.) Karst. 1881. Also pathogens of dangerous fungal diseases were found on Semyonov fir (*Trichothecium roseum* (Pers.) Link 1809 and *Myxotrichum chartarum* Kunze, 1823).

Key words: fir; fir forests; fungi; pathogens; nursery pests; fungicides; biodiversity; diseases; seedlings.

Introduction. Today the forestry of the Kyrgyz Republic is facing the needs to solve interrelated social, economic and environmental problems. Current and future forests state is alarming. A number of forest ranges in Kyrgyzstan are distressed and have lost their biological stability.

Semyonov fir or Turkestani (*A. Semonovii* Fedtsch) was first described by Professor B.A. Fedchenko in 1898 on the territory of Kyrgyzstan in the gorges of Besh-Tash of the Talas ridge. Fir trees grow in conditions of highly intersected terrain of high and medium mountains. These forests were formed on the basis of elements of tertiary complex of coniferous and coniferous-deciduous forests. They were separated from the Altai coniferous forests as a result of the Tien-Shan break. Their range is confined to the Western Tien Shan, Chatkal, Talas, At-Oynok, Uzun-Akhmat, Susamyr ranges and the Kokirim-Too Mountains. Plantings of the III-IV classes of bonitet with completeness from 0,4-0,8 predominate. They are confined mainly to the slopes of the northern exposure and occupy slopes from 1700-2700 m above sea level, where mountain-forest dark-brown soils of different thickness are common.

Semyonov fir in the past, had a common area with such species as Siberian fir, Himalayan fir and Gembila, and they may have descended from a common ancestor. Among them, only Siberian fir has the greatest geographical distribution. The main differences between it and the Tien Shan fir are the shorter needles and the absence of mechanical cells in it. Quite close to the Semyonov fir are also Central Asian firs – Himalayan and Gembila, which are distinguished by longer needles and cones.

The flora of fir forests is very diverse and is associated with the fact that this area is located at the junction between the North and South of Kyrgyzstan, and therefore you can find plants from both regions here. More than 700 species of higher flowering plants have been recorded in this area, including about 150 tree and shrub species. Dark coniferous forests on the northern slopes alternate with meadow-steppes and meadows, and light coniferous forests on the southern slopes-steppes and meadow-steppes. Below the coniferous forests grow juniper forests from juniper hemispherical and Zeravshan. Even lower, the slopes are covered with shrubs, here you can also find pistachios and almonds. In the Chatkal range, together with fir, there are Tien Shan spruce, Sivers and Kyrgyz apple trees, walnut, Turkestan and Semyonov maple, karkas, Tien Shan and Persian mountain ash, and hawthorn species (Turkestan, Dzungarian, Pontic and Altai). Along the floodplains of the rivers there are various types of willows and poplars, Turkestan birch, curve, maples, Sogdian ash and magalebka cherry. From the bushes there are various types of rosehips: loose, broad-thorned, Albert, Fedchenko, exochord Tien Shan, spirea joey leaf, Altman's and Karelin's honeysuckle, barberry oblong and variegated, Sogdian and Ferghana plum, aflatunia elm leaf, Meyer's currant, blue blackberry, cherries: Tien Shan, red-fruited, Alai and abelia [Bikirov Sh.B., 2011].

Great damage to these forests was caused by mining felling, in which only the most valuable, the best specimens of fir were selected. This led to a decrease in the genetic potential of the species, a decrease in the protective, sanitary-hygienic and aest Introduction.etic properties of the forest. In this regard, the Semyonov fir was listed in the Red Book of Kyrgyzstan as a rare endemic species.

Research methods. The research work was carried out in two stages, in the field and in the laboratory. The first stage is the allocation of permanent and temporary sample areas, and the collection of samples for mycological studies. As a result, according to the method [Zhukov, A.M.,

2001; Zhuravlev I. I., 1954], the selected samples were stored in specially sterilized bags. Growing trees were evaluated according to the four growth categories proposed by [Sinadsky Yu. V., 1977]. The degree of damage to the substrates was determined by the destruction scale [Gordienko, P. V., 1985]. At the second stage of the research, the species composition of the fungi was determined. For this purpose, preparations were used [Zhuravlev I. I., 1954] in accordance with the wet chamber method. Tinder fungi are defined according to the scheme [Komarova, E. P., 1964]. Simple microscopic preparations were prepared for fungal spores [Kirai, Z., 1974; Komarova, E. P., 1964]. Phenological and microclimatic observations in the nursery were carried out by generally accepted methods.

Research result. Semyonov fir in the Kyrgyz Republic currently occupies 3714 hectares of forests. As a result of the survey in the remote gorges of the mountains, we have identified the most valuable natural populations (genetic reserves) of the Semyonov fir, which are of great scientific, historical and economic importance. They are located in the Toktogul forestry: in the tracts of Karo-Karagay, Bugulu-Tor, Kan-Jailoo, Usta-Sai, Kuraryk, Itagar, in the Avletim forestry in the tracts of Batrakhan. The plantings here consist mainly of well-developed, healthy, multi-aged fir trees. Completeness is from 0.5 to 1.0. The optimum growth of Semyonov fir is confined to altitudes-2000-2500 m above sea level, the height of individual specimens reaches more than 30 m with a diameter of more than 1 m. The Semyonov fir tree lives up to 300 years.

Studies have found that the greatest harm to fir is caused by pests of cones, berries and seeds. As a result of the research, it was revealed that fir cones and seeds are damaged by the fir cone firefly (*Dioryctria abietella* Schiff.) and seed beetles of the genus (*Megastigmus Dalm.*).

In the years of good harvests, the spruce cone firefly of the Lepidoptera order damages a single number of fir cones up to 2.2 %, and in a low-yielding year, the activity of the firefly increases, and it damages up to 17% of the cones [Bikirov Sh. B., 1984]. This pest [Ashimov K. S., Kachibekova E. N., 1998] damages up to 90% of cones. Infected bumps are twisted. The caterpillars feed on scales, seeds, strongly wear out the cones, gnaw the rod. The biology of the pest has not been sufficiently studied. The larvae feed in cones until the end of September, then they overwinter in cocoons, which are white, soft, translucent covers with which the caterpillar covers its body, attaching to the scales of the cone.

A characteristic feature of the cones and seeds pests is a hidden lifestyle during the feeding period. The main harm they cause in the larval stage. Pests of cones and fir seeds negatively affect the volume and quality of the annual crop, destroying most of it or the entire crop. All this worsens the sanitary condition of forests and their natural renewal, and also leads to an undesirable replacement of valuable forest-forming species by less valuable in economical means. Damaged seeds, collected together with undamaged ones, reduce the grade of the harvested seeds. The most dangerous, among all above-listed pests of fir seeds is the fir seed eater. In fact, fir forests in some cases are infection foci of seed eaters of *Megastigmus Dalm.* Usually, the infestation of fir seeds with seed eaters annually amounts to 25-30 %, in some years it reaches 90%. In production conditions, large areas occupied by fir forests, the protection of fir cones from fir seed eaters is not carried out, except for special chemical control measures, in the form of various experimental developments carried out during scientific research on a limited area of the forest, i.e. on groups of trees.

Another pest particularly seen on fir is spruce-fir brown hermes (*Aphrastasia pectinatae* Chol.) - coniferous plants pest, affecting fir, larch, spruce, various types of pines. Brown spruce-

fir hermes develops on spruce and fir. During the feeding process of the brown spruce-fir hermes, the larvae concentrate on the underside of the fir needles. With mass population, the trees are covered with abundant secretions, on which soot fungi develop. During the season, 3-4 generations of hermes can develop. Aphids feed on the lower side of the needles, and yellow spots appear on the upper side in the places of sucking. With a large number of hermes, the needles turn yellow and crumble, weakened trees die within a few years. Mature trees under normal soil conditions, optimal humidity and light are more stable, and hermes develops sporadically. The pest feeds and develops normally only on fir trees.

Control measures are preventive spraying of plants in the spring, repeatedly - during the regrowth of young shoots and with a large number of pests in the summer with one of the following chemicals: fufanon, aktellik, decis Profi, kemifos, molniya, komandor, iskra, inta-vir.

In addition to the above mentioned, pests and diseases of other tree species growing here are found on the fir forest enterprise territories of the Western Tien Shan. The main ones are leaf beetles and stem pests (barbels, bark beetles) that affect poplar, willow and elm tree species. Ringed silkworm-shrubs (mountain cherry, rosehip, etc.), Dutch disease – elm and others.

Leaf beetles and barbels cause significant damage to forestry. In some years, leaf beetles in nurseries destroy up to 100% of transplant sections with rooted cuttings of poplar and willow. Measures to control leaf beetles in nurseries, due to poor-quality chemicals obtained in plant protection services, do not give positive results. Over the past 10-15 years, the large Ferghana barbel has become widespread, which has increased its number from the valley zone and settled in the mountain zone. It damages almost all types of trees. The main reason for the rapid spread of the barbel is the large trees untimely cutting and tree surgery. In most cases, the time of carrying out these works coincides with the time of laying eggs by the barbel. As a result, on moistened fresh sections of trees, larvae emerge from the laid clutches of the barbel, which, feeding on wood, develop and completely damage the trunk part of the tree. As a result, the tree completely dies or breaks. There are no effective measures to control the barbel today [Toktoraliev B. A., et al., 2006].

Among the diseases of the Semyonov fir, the most widespread in the nursery is fusarium damage to seedlings, and in natural plantings – rust and cenangial cancer.

Planting material diseases grown in forest nurseries cause great harm to forestry, as in some cases they lead to the death of a significant part of fir seedlings, up to 30-70 %. A special danger for seedlings is fusarium, which causes lodging of plants when they are grown in a nursery. This disease causes rotting of seeds and seedlings, wilting of seedlings and seedlings. The disease is focal in nature, and the plants die in clump. The main method of fir seedlings protection from fusariosis is considered to be the active method, i.e. suppression of the causative agent of the disease at the activation time by chemicals in combination with high agricultural technics. Except available control measures against this disease, usually it is practiced only fir seeds, sprouts and seedlings treatment by a solution of potassium permanganate, which give a relatively high effect.

The fungus *Cenangium abietis* (Pers.) Rehm. it was identified in the South of Kyrgyzstan as the causative agent of the dangerous fir disease "cenangial cancer", which leads dry out of the shoots in the affected areas, as well as yellowing and falling of needles. In the end, the disease leads to the death of the entire plant, the undergrowth especially suffers. It should be noted that this disease appeared relatively recently in the areas where Semyonov fir grows and has already caused significant damage in the Sary-Chelek State Biosphere Reserve [Totubaeva N. E., et al., 2003]. As one of the methods of biological control of coenangial cancer of fir, it is recommended

to control harmful insects (goldfish, sawfly, bark beetle) that damage both young and old trees and thereby contribute to the penetration of infection into plants. Also, as an additional control measure, it is recommended to select or introduce tree forms that are resistant to the most dangerous diseases [Ashimov K. S., 2004; Bilder I. V., 1999].

In natural forest stands, Semyonov fir is most strongly affected by rust fungi, they affect the branchwoods of fir trees. A characteristic sign of the affection is the thickening and cracking of the branchwoods in the affected area. In the spring, spore clumps appear on the affected branches in the form of a gelatinous yellow-brownish mass. Spores develop in the summer on an intermediate host (hawthorn, junberry, etc.), falling on the fir branches, germinate and give a long-term mycelium, which causes thickening and curvature of the branches. Usually rust begins to affect the lower branches of all ages fir at different heights and exposures, but most often in the lower and middle mountains on the northern slopes. The damage caused by rust is significant, as the disease weakens the plants, and infected branches wither. The rust (*Melampsorella cerastii* Wint.) found on the fir forms witch-brooms and tumors on the branchwoods [Mosolova S. N., Prikhodko S. L., 2006].

In addition, rust-weakened trees are subject to colonization by stem pest species, such as bark beetles, barbels, and goldfinches. Rust control measures are not carried out in fir forests, due to the high cost of work.

Another most dangerous disease of Semyonov fir can be called root rot, caused by the pine fungus *Fomitopsis annosa* (Fr.) Karst. The pine fungus is able to spread through the roots to nearby growing trees, causing a lesion focus. Infection of trees occurs with spores or mycelium, in places where the roots or lenticells of healthy roots are injured. Then the mycelium penetrates the trunk, thereby causing central rot. The bioecology of this disease was thoroughly studied by A. M. Soloviev [Soloviev A.M., 1964]. The author found that the main and permanent reservation of fungus spores is the forest litter, where more than 60% of spores are retained. Infestation of trees most often occurs by spores penetrating into the cracks of the bark and on the surface roots, or by mycelium of the forest litter. Pine fungus affects plants older than 2-3 years of age. The infestation increases with the trees age. The thickest trees are affected more strongly and reach 35-45%, all affected trees have destructive wood or die on the root. Infection of Semyonov fir with pine fungus at the age of 60-100

years reaches from 25 to 40%, and at the age of 100-120 years up to 50% of stands [Soloviev A.M., 1964]. As control measures, the following measures are recommended: cleaning, cutting, pruning and burning of dead-wood, chat wood, affected trees or individual branchwoods of needles. Decontaminate fresh wounds or other injuries with creosote oil, 5% solution of copper sulfate, 3% solution of sodium fluoride, or during the growing season 3-4 times with 1% solution of Bordeaux liquid.

Another most common disease – is alternaria blight, caused by an imperfecti fungi of *Alternaria* species, in terms of its biology, ecology and the nature of the damage caused by this fungi, is close to the representatives of *Fusarium* species, often codominant them. The *Alternaria alternate* (Fr.) Keissler species has been recorded on the Semyonov fir. It mainly affects young needles on 1-3 year-old sprouts, on trees of different ages [Sagitov A. O., et al., 2008]. The initial stage of the disease is needles yellowing, then the tip of the sprouts dies, and a black plaque forms on the dead needles, which is a septic (multi-layered, compacted) thick-walled fungi mycelium.

Disease signs on the seedlings are noticeable in the spring, in autumn they turn yellow, and die by the spring of the next year.

In addition, on the seed skin of the Semyonov fir, pathogens of hazardous fungal diseases (*Trichothecium roseum* and *Myxotrichum chartarum*) were found, which destroy from 20 to 40% of the seeds within 7 days with sufficient humidity [Kalykova G. N., 2009].

Conclusion:

1. In general, the existence of entomological and phytopathological pests in the forest is also a part of the entire forest biodiversity created by nature itself. The sanitary condition of fir forests can be assumed relatively positive. However, hazardous organisms exist there. In the process of vital activity of these hazardous organisms, healthy trees dry out, plantings lose their role and ability to natural renewal. In order to prevent the focus of the disease and localization, it is necessary to carry out the following measures: Organize continuous monitoring of fir forests. Conduct comprehensive research activities aimed at preventing the disease, selection of the persistent forms and propagation;

2. Prohibit temporary and permanent cattle camp and grazing of livestock in the forest, especially in the floodplain part of it;

3. Clean the forest and cutting areas from felling residues, windthrows and windbreaks and dry-topped dead fir trees. Trim and remove the lower wilted or shriveled branches of the affected mature tree. Clean up the affected young stands (up to 20 cm in diameter at stump height) and the dead fir undergrowth, collect it in a specially designated place and eliminate it by burning;

4. Strengthen the fir stands protection, conduct regular monitoring, consider quarantine declaring in the focus of diseases.

References:

1. **Zhuravlev, I.I.** Diagnosis of diseases of tree species by signs accessible to the naked eye [Text] / I.I. Zhuravlev. - L.: Publishing house of the Central Scientific Research Institute of Forestry, 1954. - p.76. (in Eng.).
2. **Komarova, E.P.** Determinant of the polypores of Belarus [Text] / E.P. Komarova - Minsk: Science and Technology, 1964. - p.343. (in Eng.).
3. **Soloviev, A.M.** Pine fungus in the fir forests of Kazakhstan Altai. - Alma-Ata, 1964. - P. 132-135. (in Eng.).
4. **Kirai, Z.** Methods of Phytopathology [Text] / [Z. Kiray, Z. Klement, F. Shoimoshi, J. Veresh]. - M.: Kolos, 1974. - p. 344. (in Eng.).
5. **Sinadsky, Yu. V.** Course of lectures on forest phytopathology [Text] / Yu. V. Sinadsky - M.: Publishing House of Moscow University, 1977. - p.214. (in Eng.).
6. **Bikirov, Sh.B.** Fir forests of Kyrgyzstan [Text] / Sh.B. Bikirov. - F.: Ilim, 1984. - p.148. (in Eng.).
7. **Gordienko, P.V.** Peculiarities of settlement of wood-destroying fungi on a substrate of varying degrees of destruction in the mixed coniferous-broad leaved forest of Sikhote-Alin. [Text] / P.V. Gordienko // Fungi and lichens in the ecosystem // Abstract report of the X scientific symposium of mycologists and lichenologists of the Baltic republics and Belarus. Part 1. - Riga: Leningrad State University, 1985. - P. 44-46. (in Eng.).
8. **Ashimov, K.S.** Dendrophilous insects and diseases of mountain forests in the South of Kyrgyzstan [Text] / K.S. Ashimov, E.N. Kachibekova, M.Zh.Nurmanbaev // Bioecology of walnut forests and geodynamics in Southern Kyrgyzstan. Publication II. - Zhalal-Abad, 1998. - P. 107-113. (in Eng.).
9. **Bilder, I.V.** Methods of combating fungal diseases of spruce and fir in Kyrgyzstan [Text] / I.V. Bilder // Recommendations on forestry issues in Kyrgyzstan. - Bishkek: Niva, 1999. - P. 82-89. (in Eng.).

10. **Zhukov, A.M.** Scientific-methodical manual for the diagnosis of fungal diseases of flax trees and shrubs [Text] / A.M. Zhukov, P.V. Gordienko. - M.: Publishing house of the Ministry of Natural Resources of Russia, 2001. - p. 69. (in Eng.).
11. **Totubaeva, N.E.** Sanitary condition of Semyonov fir in the buffer zone of Sary-Chelek nature reserve [Text] / [N.E. Totubaeva, Sh.B. Bikirov, T.O. Zhunusov, S.Kenzhebaev] // Silvicultural research in Kyrgyzstan, Bishkek, 2003.- P. 161-167. (in Eng.).
12. **Ashimov, K.S.** Melampsorella cerastii of Semyonov fir. [Text] / K.S. Ashimov // Study of wildlife of Kyrgyzstan. - Bishkek, 2004. – Publication 5.- P. 196-197. (in Eng.).
13. **Ashimov, K.S.** Fungal diseases of Semyonov fir in the Chichkan gorge of Toktogul forestry [Text] / K.S. Ashimov, G.N. Kalykova // Bulletin of Zhalal-Abad State University. - Zhalal-Abad, 2006. - p. 67-70. (in Eng.).
14. **Mosolova, S.N.** State of mycological studies of the main forest-forming species in Kyrgyzstan [Text] / S.N. Mosolova, S.L. Prikhodko // Rational use and conservation of forest resources. – Publication 21. Bishkek, 2006. - P. 198-203. (in Eng.).
15. **Toktoraliyev, B.A.** Guide to growing juniper seedlings in nurseries, protecting them from diseases and pests [Text] / [B.A. Toktoraliyev, A.V. Kosmynin, Sh.B. Bikirov, A.T. Attokurov] // Publishing house of OshTU.- Osh: TU, 2006. - p.31. (in Eng.).
16. **Sagitov, A.O.** Alternaria blight of Semenovii fir in Kyrgyzstan. Bulletin of the National Academy of Sciences of the Kyrgyz Republic [Text] / A.O. Sagitov, A.S. Kochorov, G.N. Kalykova. – B.: Ilim, 2008.- No.3.- P.100-102. (in Eng.).
17. **Sagitov, A.O.** Pine fungus of Semenovii fir in the Western Tien Shan [Text] / A.O. Sagitov, K.S. Ashimov, G.N. Kalykova // Actual problems of forest management and staffing of the forestry sector of the economy of Central Asia. - International Scientific and Practical Conference.- Almaty Agrouniversity Publishing House, 2008.- P. 254-257. (in Eng.).
18. **Kalykova, G.N.** Diseases of white pine seeds and the fight against them [Text] / G.N. Kalykova // Bulletin of Kyrgyz University of Agriculture.- No. 4 (15).- Bishkek, 2009. – P. 117-120. (in Eng.).
19. **Bikirov, Sh.B.** Scientific basis for conservation and restoration of fir forests in the Western Tien Shan [Text] / Sh.B. Bikirov.-Bishkek: Polygraphbumresursy, 2011 – p. 396. (in Eng.).

DOI: 10. 54834 / 16945220_2021_3_66

Поступила в редакцию 19. 09. 2021 г.